

The σ meson in a nuclear medium through two pion photoproduction

L. ROCA, E. OSET AND M. J. VICENTE VACAS

Depto. de Física Teórica and IFIC. Centro Mixto Universidad de Valencia-CSIC

We present theoretical results for $(\gamma, \pi^0\pi^0)$ and $(\gamma, \pi^\pm\pi^0)$ production on nucleons and nuclei in the kinematical region where the scalar isoscalar $\pi\pi$ amplitude is influenced by the σ pole. The final state interaction of the pions modified by the nuclear medium produces a spectacular shift of strength of the $\pi^0\pi^0$ invariant mass distribution induced by the moving of the σ pole to lower masses and widths as the nuclear density increases.

PACS numbers: 25.20.Lj, 14.40.Cs, 12.39.Fe, 21.30.Fe

In the last years there has been an intense theoretical and experimental debate about the nature of the σ meson, especially the modification of its properties in nuclear matter. Both the interpretation as a $q\bar{q}$ state or as a $\pi\pi$ resonance, seem to agree in a decrease of the σ mass as the nuclear density increases. For instance, in the theoretical work of [3], the $\pi\pi$ amplitude in the σ channel showed a shift of strength towards low $\pi\pi$ invariant masses at finite nuclear density. In the present work, our purpose is to find the possible experimental signature of this moving of the $\pi\pi$ pole in the $(\gamma, \pi^0\pi^0)$ reaction in nuclei. This reaction is much better suited than the $(\pi, \pi\pi)$ one to investigate the modification of the $\pi\pi$ in nuclear matter because the photons are not distorted by the nucleus and one can test higher densities.

For the model of the elementary $(\gamma, \pi\pi)$ reaction we follow [2] which considers the coupling of the photons to mesons, nucleons, and the resonances $\Delta(1232)$, $N^*(1440)$, $N^*(1520)$ and $\Delta(1700)$. This model relies upon tree level diagrams. Final state interaction of the πN system is accounted for by means of the explicit use of resonances with their widths. However, since we do not include explicitly the σ resonance, the final state interaction of the two pions has to be implemented to generate it.

The $\gamma N \rightarrow N\pi^0\pi^0$ amplitude can be decomposed in a part which has in the final state the combination of pions in isospin $I=0$ and another part where the pions are in $I=2$.

The renormalization of the $I = 0$ ($\gamma, \pi\pi$) amplitude is done by factorizing the on shell tree level $\pi N \rightarrow \pi\pi N$ and $\pi\pi \rightarrow \pi\pi$ amplitudes in the loop functions.

$$T_{(\gamma, \pi^0 \pi^0)}^{I_{\pi\pi}=0} \rightarrow T_{(\gamma, \pi^0 \pi^0)}^{I_{\pi\pi}=0} \left(1 + G_{\pi\pi} t_{\pi\pi}^{I=0}(M_I) \right) \quad (1)$$

where $G_{\pi\pi}$ is the loop function of the two pion propagators, which appears in the Bethe Salpeter equation, and $t_{\pi\pi}^{I=0}$ is the $\pi\pi$ scattering matrix in isospin $I=0$, taken from [3]. In the model for $(\gamma, 2\pi)$ of [2] there are indeed contact terms as implied before, as well as other terms involving intermediate nucleon states or resonances. In this latter case the loop function involves three propagators but the intermediate baryon is far off shell and the factorization of Eq. (1) still holds. There is, however, an exception in the Δ Kroll Ruderman term, since as we increase the photon energy we get closer to the Δ pole. For this reason this term has been dealt separately making the explicit calculation of the loop with one Δ and two pion propagators.

The cross section for the process in nuclei is calculated using many body techniques. The distortion of the final pions in their way out through the nucleus via absorption and quasielastic collisions is of crucial importance and is taken into account by using an eikonal approximation. The probability of absorption and quasielastic collisions is calculated using a well tested model for the pion selfenergy in nuclear matter for low energy pions. With this approximation the pions which undergo absorptions are removed from the flux but we do not remove those which undergo quasielastic collisions since they do not change in average the shape or the strength of the $\pi\pi$ invariant mass distribution.

In the figure we can see the results for the two pion invariant mass distributions in the $(\gamma, \pi^0 \pi^0)$ and $(\gamma, \pi^\pm \pi^0)$ reactions on ^{12}C and ^{208}Pb . The difference between the solid and dashed curves is the use of the in medium $\pi\pi$ scattering and G function instead of the free ones, which we take from [3]. As one can see in the figure, there is an appreciable shift of strength to the low invariant mass region in the $(\gamma, \pi^0 \pi^0)$ due to the in medium $\pi\pi$ interaction. This shift is remarkably similar to the one found in the measurements of [4]. This shift is not seen in the $(\gamma, \pi^\pm \pi^0)$ channel because the $\pi^\pm \pi^0$ are not allowed to be in isospin $I=0$.

These results show a clear signature of the modified $\pi\pi$ interaction in the nuclear medium that can be related to the modification of the σ meson in the medium since, in the model we use, the σ meson is dynamically generated by the multiple scattering of the pions through the underlying chiral dynamics.




Fig. 1. Two pion invariant mass distribution for $\pi^0\pi^0$ and $\pi^\pm\pi^0$ photoproduction in ^{12}C and ^{208}Pb . Continuous lines: using the in medium final $\pi\pi$ interaction. Dashed lines: using the final $\pi\pi$ interaction at zero density. Exp. data from [4]

REFERENCES

- [1] L. Roca, E. Oset and M. J. Vicente Vacas, arXiv:nucl-th/0201054.
- [2] J. C. Nacher, E. Oset, M. J. Vicente and L. Roca, Nucl. Phys. A **695** (2001) 295 [arXiv:nucl-th/0012065].
- [3] H. C. Chiang, E. Oset and M. J. Vicente-Vacas, Nucl. Phys. A **644** (1998) 77 [arXiv:nucl-th/9712047].
- [4] J. G. Messchendorp *et al.*, arXiv:nucl-ex/0205009.